

Présentation du cadre HSPA pour le Luxembourg



Lucie Bryndová
OCDE



Till Seuring
ObSanté



Anne-Charlotte Lorcy
ObSanté



Kadri-Ann Kallas
OCDE



OBSERVATOIRE
NATIONAL DE LA **SANTÉ**



OCDE
DES POLITIQUES MEILLEURES
POUR UNE VIE MEILLEURE



Financé par
l'Union européenne

HSPA use case study: Mental Health



Kadri-Ann Kallas
OCDE



OBSERVATOIRE
NATIONAL DE LA SANTÉ



OCDE
DES POLITIQUES MEILLEURES
POUR UNE VIE MEILLEURE




Financé par
l'Union européenne



Mental health as an HSPA use case


- > Policy-relevant and priority in Luxembourg
- > Visible in the framework
 - > dedicated mental health subdomain





Mental health as an HSPA use case


- Policy-relevant and priority in Luxembourg
- Visible in the framework
 - dedicated mental health subdomain
 - But also reflected across the framework: indicators in multiple domains






Inputs: capacity to deliver care

Domain and subdomain	Name of the indicator	Indicator breakdown	HSPA indicator custodian
Infrastructure/ Physical resources	Hospital beds	Ownership status, healthcare function, care type	ObSanté
Workforce/ Workforce capacity	Practicing physicians	Age, sex, specialisation, country of birth, country of residence	ObSanté





Psychiatric hospital beds per 1000 population



Processes: care organisation and delivery

Domain and subdomain	Name of the indicator	Indicator breakdown	HSPA indicator custodian
Access to care/ Financial protection	Unmet need for mental healthcare due to financial reasons	Sex, age, income quintile and level of disability	ObSsanté
Value-based care/ Cost-efficiency	In-patient average length of stay	Age, sex, cause of disease (ICD-10) ²	ObSanté
Integrated care/ Co-ordination of care	Experienced co-ordination: patients with chronic conditions who report having a care plan available to them	Age, gender, education and more	ObSanté
Integrated care/ Co-ordination of care	Experienced co-ordination: patients with chronic conditions who report professionals in their care appear to be following the same care plan	Age, gender, education and more	ObSanté
Integrated care/ Continuity of care	Suicide within 1 year after discharge among patients diagnosed with a mental disorder		ObSanté







Outcomes: mental health status



Domain and subdomain	Name of the indicator	Indicator breakdown	HSPA indicator custodian
Health status/ Mental health	Proportion (%) of adolescents reporting multiple health complaints	Gender, perceived wealth (affluence)	ObSanté
Health status/ Mental health	Current depressive symptoms	Sex, age, country of birth	ObSanté
Health status/ Mental health	Emotional distress (feeling anxious, depressed or irritable)	Age, gender, income	
Health status/ Population outcomes	Distribution of beneficiaries of LTC	Morbidity, disability (e.g. hearing)	
Person-centredness/ Health literacy	Mental health literacy		




Panel A. 15-year-olds who report multiple health complaints more than once a week






Outcomes: mental health status

Domain and subdomain	Name of the indicator	Indicator breakdown	HSPA indicator custodian
Health status/ Mental health	Proportion (%) of adolescents reporting multiple health complaints	Gender, perceived wealth (affluence)	ObSanté
Health status/ Mental health	Current depressive symptoms	Sex, age, country of birth	ObSanté
Health status/ Mental health	Emotional distress (feeling anxious, depressed or irritable)	Age, education	
Health status/ Population outcomes	Distribution of beneficiaries of LTC	Morbidity (e.g. health)	
Person-centredness/ Health literacy	Mental health literacy		





Panel B. Current depressive symptoms among adults





Cross-cutting: influence care provision and outcomes

Domain and subdomain	Name of the indicator	Indicator breakdown	HSPA indicator custodian
Determinants of health/ Behavioural risk factors	Frequency of alcohol consumption	Sex, age, education, income	ObSanté






HSPA indicators in a pathway: Mental Health

- > **Determinants** (alcohol)
- > **Capacity** (beds, psychiatrists)
- > **Access & delivery** (financial unmet need, length of stays, coordination)
- > **Population outcomes** (adolescents, depressive symptoms, distress, LTC)

- > This case study highlights:
 - > Connections between (sub)domains
 - > Data is already available for many indicators
 - > Possible illustrations

Moving beyond isolated indicators towards a more coherent understanding of performance



 Francesca.colombo@oecd.org

lucie.bryndova@oecd.org

kadri-ann.kallas@oecd.org



 hspa@obs.etat.lu

Thank you.

 www.oecd.org/health/

 <https://www.oecd.org/health/health-system-performance-assessment.htm/>

French version will be available in May 2026



OBSERVATOIRE
NATIONAL DE LA SANTÉ



OCDE

DES POLITIQUES MEILLEURES
POUR UNE VIE MEILLEURE



Financé par
l'Union européenne